



ROR2 gene

receptor tyrosine kinase like orphan receptor 2

Normal Function

The *ROR2* gene provides instructions for making a protein whose function is not well understood. The ROR2 protein is part of a family of proteins known as receptor tyrosine kinases (RTKs), which play a role in chemical signaling within cells. RTKs are involved in many cell functions, including cell growth and division (proliferation), the process by which cells mature to carry out specific functions (differentiation), cell survival, and cell movement (motility).

Researchers believe that the ROR2 protein plays an essential role in development starting before birth. It is involved in chemical signaling pathways called Wnt signaling, which affect many aspects of development. These pathways control the activity of genes needed at specific times, and they regulate the interactions between cells as organs and tissues are forming. In particular, the ROR2 protein appears to be critical for the normal formation of the skeleton, heart, and genitals.

Health Conditions Related to Genetic Changes

Robinow syndrome

At least 16 mutations in the *ROR2* gene have been found to cause the autosomal recessive form of Robinow syndrome, a condition that affects the development of many parts of the body, particularly the bones. Autosomal recessive inheritance means both copies of the gene in each cell have mutations. Some of these mutations change single protein building blocks (amino acids) in the ROR2 protein, while others lead to the production of an abnormally short, nonfunctional version of the protein. Because these genetic changes prevent any functional ROR2 protein from being made, they are described as "loss-of-function" mutations. Loss of the ROR2 protein impairs chemical signaling pathways that are important for normal development, particularly the formation of bones in the face, spine, and limbs. These changes lead to the skeletal abnormalities characteristic of Robinow syndrome. A lack of this protein during early development also underlies the other features of Robinow syndrome, including genital abnormalities and heart defects.

other disorders


More than 10 mutations in the *ROR2* gene have been identified in people with a disorder called brachydactyly type B1. This condition is characterized by abnormally short fingers and toes, particularly the fourth and fifth digits, and malformed or

absent fingernails and toenails. Brachydactyly type B1 has an autosomal dominant pattern of inheritance, which means one altered copy of the gene in each cell is sufficient to cause the disorder. Unlike the mutations that cause Robinow syndrome (described above), the *ROR2* gene mutations responsible for brachydactyly type B1 are described as "gain-of-function" because they appear to cause the ROR2 protein to be continuously active. It is unclear how the overactive protein disrupts the development of bones in the hands and feet.

Chromosomal Location

Cytogenetic Location: 9q22.31, which is the long (q) arm of chromosome 9 at position 22.31

Molecular Location: base pairs 91,722,596 to 91,950,206 on chromosome 9 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- BDB1
- brachydactyly type B1 gene
- neurotrophic tyrosine kinase receptor-related 2 gene
- NTRKR2
- receptor tyrosine kinase-like orphan receptor 2
- ROR2_HUMAN

Additional Information & Resources

GeneReviews

- ROR2-Related Robinow Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1240>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28ROR2%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- RECEPTOR TYROSINE KINASE-LIKE ORPHAN RECEPTOR 2
<http://omim.org/entry/602337>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
<http://atlasgeneticsoncology.org/Genes/ROR2ID43476ch9q22.html>
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=ROR2%5Bgene%5D>
- HGNC Gene Family: I-set domain containing
<http://www.genenames.org/cgi-bin/genefamilies/set/593>
- HGNC Gene Family: Receptor Tyrosine Kinases
<http://www.genenames.org/cgi-bin/genefamilies/set/321>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=10257
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/4920>
- UniProt
<http://www.uniprot.org/uniprot/Q01974>

Sources for This Summary

- Afzal AR, Jeffery S. One gene, two phenotypes: ROR2 mutations in autosomal recessive Robinow syndrome and autosomal dominant brachydactyly type B. *Hum Mutat.* 2003 Jul;22(1):1-11.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12815588>
- Afzal AR, Rajab A, Fenske CD, Oldridge M, Elanko N, Ternes-Pereira E, Tüysüz B, Murday VA, Patton MA, Wilkie AO, Jeffery S. Recessive Robinow syndrome, allelic to dominant brachydactyly type B, is caused by mutation of ROR2. *Nat Genet.* 2000 Aug;25(4):419-22.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10932186>
- GeneReview: ROR2-Related Robinow Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1240>

- Oldridge M, Fortuna AM, Maringa M, Propping P, Mansour S, Pollitt C, DeChiara TM, Kimble RB, Valenzuela DM, Yancopoulos GD, Wilkie AO. Dominant mutations in ROR2, encoding an orphan receptor tyrosine kinase, cause brachydactyly type B. *Nat Genet.* 2000 Mar;24(3):275-8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10700182>
- Schwabe GC, Tinschert S, Buschow C, Meinecke P, Wolff G, Gillessen-Kaesbach G, Oldridge M, Wilkie AO, Kömec R, Mundlos S. Distinct mutations in the receptor tyrosine kinase gene ROR2 cause brachydactyly type B. *Am J Hum Genet.* 2000 Oct;67(4):822-31. Epub 2000 Sep 12.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10986040>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1287887/>
- Stricker S, Mundlos S. FGF and ROR2 receptor tyrosine kinase signaling in human skeletal development. *Curr Top Dev Biol.* 2011;97:179-206. doi: 10.1016/B978-0-12-385975-4.00013-9. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22074606>
- Wang B, Sinha T, Jiao K, Serra R, Wang J. Disruption of PCP signaling causes limb morphogenesis and skeletal defects and may underlie Robinow syndrome and brachydactyly type B. *Hum Mol Genet.* 2011 Jan 15;20(2):271-85. doi: 10.1093/hmg/ddq462. Epub 2010 Oct 20.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20962035>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3031336/>
- van Bokhoven H, Celli J, Kayserili H, van Beusekom E, Balci S, Brussel W, Skovby F, Kerr B, Percin EF, Akarsu N, Brunner HG. Mutation of the gene encoding the ROR2 tyrosine kinase causes autosomal recessive Robinow syndrome. *Nat Genet.* 2000 Aug;25(4):423-6.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10932187>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/ROR2>

Reviewed: September 2015
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services